

RESEARCH PAPER

Morphometrics and biological characteristics of *Pentalonia nigronervosa*, the vector of *Banana bunchy top virus*, living on various Araceous plants species

Suparman Suparman¹, Rafika Oktarida², Harman Hamidson¹, & Arsi Arsi¹

Manuscript received: 19 November 2022. Revision accepted: 23 January 2023. Available online: 3 March 2023.

ABSTRACT

Banana aphid *Pentalonia nigronervosa* Coquerel (Hemiptera: Aphididae) is widely known as the main vector of *Banana bunchy top virus* (BBTV). It had been reported that banana aphid could live on alternative hosts belong to Family Araceae, and become important issue since Araceous plants are commonly found surrounding banana cultivations. Research had been conducted to study the morphometrics and biological characteristics of banana aphids living on various species of Araceous plants. The results showed that Araceous plant species could influence the morphometric and biological characteristic of banana aphids, indicated by the significant difference in some biological characteristics such as life cycle, life span, reproductive period, and fecundity. The average life span of banana aphids living on Araceous plants ranged from 22.80 to 47.90 days, it was shorter than the one that of lived on a banana plants. *P. nigronervosa* was able to develop a bigger colony when the aphid lived on Araceous plants, especially when they were transferred to Bogor taro where their fecundity reached 63 nymphs per female. Araceous plants also affected the morphometric of *P. nigronervosa* indicated by smaller size of the aphids, compared to those living on banana plant. The ability of banana aphid to live and reproduce in various Araceous plants would be beneficial to the development of control strategies of BBTV.

Key words: banana aphid, life cycle, life span, fecundity

INTRODUCTION

Banana aphid *Pentalonia nigronervosa* Coquerel (Hemiptera: Aphididae) is a major pest of banana (Biale et al., 2017) and is the main vector of *Banana bunchy top virus* (BBTV). The role of the aphid as the vector of BBTV make the aphid more dangerous to banana cultivation (Watanabe et al., 2013). The virus is considered to be the most important pathogen of bananas which might cause yield losses up to 100% (Qazi, 2016). Infected banana produces no fruit, but if the plant is infected at later stage, small fruits might be produced but not marketable (Jekayinoluwa et al., 2020). The first symptom of BBTV infection is the appearance of dark green broken lines on infected plant leaves (Chen & Hu, 2013). At the later stage of infection, newly formed leaves are shorter and narrower with chlorotic areas appear on leaf margins (Elayabalan et al., 2015).

P. nigronervosa generally lives in association with banana plants worldwide, including tropical and subtropical regions (Footitt & Maw, 2019). Banana aphids prefer to stay on the lower part of the host plant where they can build an association with ants, from which they get protection from predators (Biale et al., 2017). Instead of being the major pest of banana, *P. nigronervosa* could also live on various alternative hosts belonging to Families Araceae and Zingiberaceae (Suparman et al., 2017).

P. nigronervosa transmit BBTV in a circulative manner through the feeding activity (Niyongere et al., 2013; Jebakumar et al., 2018; Jekayinoluwa et al., 2021). The virus particles enter the aphid body together with plant liquid sucked by the aphid from the phloem of infected banana. The virus particles move from foregut to midgut and hemocoel before reaching salivary gland, from where the virus particles are transmitted to healthy plants through the aphid stylet (Watanabe & Bressan, 2013). BBTV particles were acquired by *P. nigronervosa* after an acquisition feeding period of 15 min and a single aphid was able to acquire 861 virus particles after 24 h of feeding period (Jebakumar et al., 2018). In the transmission of the virus, the vector is influenced by many factors such as host plant conditions, temperature, and the life stage of the vector itself (Bragard et al., 2013). Environmental conditions,

Corresponding author:

Suparman Suparman (suparmanshk@gmail.com)

¹Department of Plant Protection, Faculty of Agriculture, Universitas Sriwijaya. Jl. Raya Palembang-Prabumulih Km 32, Indralaya, Ogan Ilir, South Sumatra, Indonesia 30662

²Crop Sciences Graduate Program, Faculty of Agriculture, Universitas Sriwijaya. Jl. Padang Selasa No. 524, Bukit Besar, Palembang, South Sumatra, Indonesia 30139

especially temperature, could affect the population of *P. nigronervosa* due to parthenogenetic reproduction of the aphid which reached the highest rate of population development at the temperature of 25 °C. At the same temperature, the life cycle of *P. nigronervosa* ranged from 10 to 14 days and reproduction period ranged from 9 to 12 days with fecundity 20 to 34 nymphs per female (Basak et al., 2015).

Host species could influence morphological and physiological change of aphids (Mdellel & Kamel, 2015). Different host species could provide different food and environment for aphid colonization (Le Guigo et al., 2012), and *P. nigronervosa* showed different reproduction capacities when lived on different alternative host species (Suparman et al., 2017).

BBTV is very difficult to control due to the virus itself and the vector involved. The virus is able to persist for quite long time in cut infected banana pseudo stem and still transmittable to healthy banana (Tricahyati et al., 2022), and persists for even longer in the infected corm. The vector has many alternative hosts and shows different performance on different host species and some host species are able to reduce transmission efficiency of the virus (Oktarida et al., 2022). Therefore, understanding the morphometric and biological characteristics of *P. nigronervosa* on each alternative host might be valuable in the design of control measures of the vector and the disease.

MATERIALS AND METHODS

Research Site. The research was conducted in the Laboratory of Entomology and experimental garden, Department of Plant Pests and Diseases, Faculty of Agriculture, Universitas Sriwijaya, Indonesia from May to December 2021.

Banana Aphid Isolation and Rearing. Banana aphid, *P. nigronervosa* was taken from banana plant belong to farmers in Ogan Ilir Regency, South Sumatra. Single aphid was then bred parthenogenetically on banana sucker in a 50 × 50 × 150 cm³ wire mesh rearing cage for three weeks to produce a relatively big colony. Thirty individual were used in this rearing and the biggest colony developed was selected for further works. The colony was then further reared strictly on banana suckers to guarantee that the research used only aphids purely reproduced from single aphid originally from banana host. Numbers of nymphs and adults of the colony had also been used to experimentally transmit BBTB and successfully transmitted the virus at high efficiency. The strict isolation and rearing of the banana aphid were

aimed at preventing the mixing of *P. nigronervosa* and *P. caladii* which was morphologically and behaviorally similar to the banana aphid (Duay et al., 2014).

The banana aphids were also bred on Araceous plants prior to the conduction of the biological assay experiment, to make the aphids familiar with the alternative host where they would live and feed on during the experiment. The aphids were only being reared on the same plant species as they would be assayed during the experiment. The Araceous plant suckers used to rear the banana aphid were grown in 20-cm diameter pots covered with a plastic transparent cylinder with a cheese cloth window for aeration. The pots were placed in a shade house and the aphids were allowed to breed and produce enough imagoes for the biological assay experiment.

Araceous Plant Preparation. The experiment on the morphometric and biological characteristics of *P. nigronervosa* was conducted by using 7 species of Araceous plants and one genotype of banana. The Araceous plants were Belitung taro (*Xanthosoma sagittifolium* (L.) H.W. Schott & Endl), Bogor taro (*Colocasia esculenta* L. Schott), Pontianak taro (*C. esculenta* (L.) Schott), Japanese taro (*C. esculenta* var. *antiquorum*), Ornamental caladium (*Caladium bicolor* Vent), Wild taro (*Colocasia* sp.) and Rodent tuber (*Typhonium flagelliforme* (L.) Bl.). The Araceous plants which were commonly found surrounding banana cultivation areas and one genotype of banana was collected from farmers' fields. The banana genotype used in the experiment was the Lady Finger banana (AAA), the most widely cultivated banana cultivar in South Sumatera. All experimental plants were planted individually to produce more than 10 equal-size suckers to facilitate the experiment which needed 10 suckers of each host species for replication.

Biological Assay of *P. nigronervosa*. The experiment to obtain morphometric and biological characteristics of *P. nigronervosa* on Araceous plants was arranged in a completely randomized design with 8 treatments and 10 replications. The treatment was *P. nigronervosa* host species, comprised of 7 Araceous plant species and 1 banana genotype. Ten suckers of equal size were used for each host species and planted separately in 10-cm diameter transparent-plastic bottles filled with 5-cm high soil at the bottom. A newly born *P. nigronervosa* nymph was carefully transferred to the sucker in each bottle by using wetted tip of fine-paint brush. Each bottle was then covered with cheese cloth to facilitate air movement and protect the nymph from predators.

The observation and measurement were made daily to collect morphometric and biological data of each nymph until the nymph became imago. The morphometrics of *P. nigronervosa* such as body length and head width were measured accordingly to the method proposed by Bagariang et al. (2019). Body length was measured from the front to the cauda apex and head width was measured across the eyes. All measurement was taken for every instar of the aphid.

Number of instars was recorded based on the number of molting that occurred from hatching until the formation of the aphid imago. The instar period was measured as a period between two successive molts of a nymph. The life cycle was measured from the aphid was born until its adult gave birth for the first time. The reproductive period was a period between the first and the last time an adult gave birth to its progenies. The life span of an aphid was a period from the aphid was born until died, and it could also be calculated by summing up all instar periods and adult period. Fecundity was measured as the number of nymphs delivered by an adult female during its reproductive period. During the counting of fecundity, a newly born nymph was taken out from the bottle after being counted (Gao et al., 2021). The counting of the progenies was conducted every 24 hours.

RESULTS AND DISCUSSION

Morphometric of *Pentalonia nigronervosa*.

Morphological characteristics of *P. nigronervosa* lived on Araceous species and banana were not significantly different from each other in the first instar, they started to vary from second instar. The body length of the aphid lived on araceous plants ranged from 0.70 to 0.84 mm in the second instar, 1.04 to 1.16 mm in the third instar and 1.23 to 1.30 mm in the fourth instar (Table 1). In the

second and third instars, the average body length of the aphid in Bogor taro and rodent tuber were significantly different from that in banana. However, in the fourth instar, only the body length of the aphid lived in Belitung taro was significantly different from that in banana.

The measurement of head width of the aphid showed that the average head width of the aphid lived in some Araceous species were significantly different from the one that lived in banana (Table 2). In the second instar, the head width in Bogor taro (0.43 mm) and Japanese taro (0.45 mm) were significantly different from that in banana (0.54 mm). In the third instar, the characteristics of aphid lived in Bogor taro (0.62 mm), Pontianak taro (0.62 mm), rodent tuber (0.67 mm), and ornamental caladium (0.67 mm) were significantly different from the one that lived in banana (0.79 mm). When the aphid reached the fourth instar, only the one that lived in Bogor taro (0.79 mm) and Pontianak taro (0.84 mm) that has different characteristics from the one that lived in banana.

The measurement of the aphid antennae also showed that significant differences of the average length of the antennae among host species was not consistent from instar to instar (Table 3). In the second instar, the aphid length of the antennae in Bogor taro (0.72 mm) and rodent tuber (0.72 mm) were significantly different from that in banana (0.84 mm). In the third instar, the length of the antennae in Bogor taro (1.07 mm), Japanese taro (1.08 mm) and rodent tuber (1.07 mm) were significantly different from that in banana (1.16 mm). In the fourth instar, the length of the antennae in Belitung taro (1.28 mm) and Pontianak taro (1.28 mm) were significantly different from that in banana (1.34 mm).

P. nigronervosa is a hemimetabolous insect undergoes incomplete metamorphosis. The insect reproduces parthenogenetically and has only 2 distinct

Table 1. Body length measurements of *Pentalonia nigronervosa* lived on various host plants

Host species	Body length (mm)			
	First instar ^{ns}	Second instar	Third instar	Fourth instar
Bogor taro	0.37 ± 0.01	0.70 ± 0.02 b	1.04 ± 0.03 b	1.25 ± 0.01 ab
Belitung taro	0.37 ± 0.01	0.73 ± 0.02 ab	1.16 ± 0.02 ab	1.23 ± 0.01 b
Pontianak taro	0.36 ± 0.02	0.81 ± 0.02 a	1.12 ± 0.01 ab	1.24 ± 0.005 ab
Japanese taro	0.36 ± 0.02	0.72 ± 0.03 ab	1.06 ± 0.01 b	1.27 ± 0.01 ab
Wild taro	0.39 ± 0.01	0.72 ± 0.04 ab	1.16 ± 0.02 a	1.30 ± 0.02 a
Rodent tuber	0.38 ± 0.02	0.70 ± 0.02 b	1.04 ± 0.04 b	1.24 ± 0.03 ab
Ornamental caladium	0.38 ± 0.01	0.79 ± 0.01 ab	1.10 ± 0.03 ab	1.25 ± 0.02 ab
Lady finger banana	0.38 ± 0.01	0.84 ± 0.04 a	1.16 ± 0.01 a	1.30 ± 0.02 a

^{ns}= not significant; Numbers followed by the same letter are not significantly different according to LSD 5%.

life stages because there is no egg stage in their life cycle. Nymphs of *P. nigronervosa* are born live and they pass 4 instars before the turn into adult stage. The first instar nymph begins at born until the first molt, and the next instars are a stage period between two successive molts. *P. nigronervosa* undergoes 4 successive molts and therefore the aphid nymphs pass 4 nymph instars.

As presented in Table 1, there was no difference of the size of the first instar of *P. nigronervosa* living on different host species. The significant difference of body size between the aphid nymphs living on banana and those living on some other hosts was found in the next three instars. Bogor seemed to be the Araceous host causing significant difference on body length and head width size of the aphid compared to those living on banana, even though the difference of body length was only significant in second and third instar. The difference of body size indicated the different growth rate of the insect, which could be caused by different quality of nutrition intake (Martínez & Costamagna, 2018).

The head width of *P. nigronervosa* was measured as the representative of the aphid body width. Unlike the aphid abdomen which is elastic and getting bigger after

feeding, the head of the aphid is rigid and its size does not change after feeding. In order to obtain the maximum head width, the measurement was taken across the aphid eyes. As presented in Table 2, the head width of *P. nigronervosa* lived on Araceous species varied in every instar except in the first instar, and the variation differed from instar to instar. In Araceous species, the head width of *P. nigronervosa* tent to be smaller than that in banana. Foottit et al. (2010) reported that the head width of *P. nigronervosa* was wider when it lived on Musaceae plants than when it lived on other hosts.

P. nigronervosa that lived on Araceous plant has similar or smaller body length and head width than those of the aphid lived on banana, indicating that the plants contained appropriate nutrition suitable to the aphids, but the quality of the nutrition contained was lower than that of other hosts, especially when being compared to the nutritional value of banana, the main host of the aphid. However, there was a report that the morphological differences among aphid living on different hosts were more likely to be caused by genetical factors rather than by environmental factors, including the nutritional quality of the hosts (Aqueel et al., 2013).

Table 2. Head width measurements of *Pentalonia nigronervosa* lived on various host plants

Host species	Head width (mm)			
	First instar ^{ns}	Second instar	Third instar	Fourth instar
Bogor taro	0.21 ± 0.01	0.43 ± 0.01 b	0.62 ± 0.03 b	0.79 ± 0.03 b
Belitung taro	0.20 ± 0.01	0.47 ± 0.03 ab	0.75 ± 0.01 ab	0.87 ± 0.02 ab
Pontianak taro	0.20 ± 0.01	0.47 ± 0.02 ab	0.62 ± 0.02 b	0.84 ± 0.03 b
Japanese taro	0.19 ± 0.02	0.45 ± 0.03 b	0.74 ± 0.03 ab	0.98 ± 0.03 ab
Wild taro	0.21 ± 0.02	0.46 ± 0.03 ab	0.76 ± 0.04 a	1.01 ± 0.05 a
Rodent tuber	0.23 ± 0.02	0.38 ± 0.02 ab	0.67 ± 0.02 b	0.93 ± 0.07 ab
Ornamental caladium	0.22 ± 0.02	0.49 ± 0.02 ab	0.67 ± 0.02 b	0.97 ± 0.04 ab
Lady finger banana	0.21 ± 0.02	0.54 ± 0.04 a	0.79 ± 0.04 a	1.00 ± 0.05 a

^{ns}= not significant; Numbers followed by the same letter are not significantly different according to LSD 5%.

Table 3. Length of antenna measurements of *Pentalonia nigronervosa* lived on various host plants

Host species	Length of antenna (mm)			
	First instar ^{ns}	Second instar	Third instar	Fourth instar ^{ns}
Bogor taro	0.40 ± 0.02	0.72 ± 0.02 b	1.07 ± 0.03 b	1.31 ± 0.02
Belitung taro	0.39 ± 0.02	0.74 ± 0.02 ab	1.17 ± 0.01 a	1.28 ± 0.004
Pontianak taro	0.37 ± 0.02	0.82 ± 0.02 a	1.14 ± 0.01 ab	1.28 ± 0.01
Japanese taro	0.38 ± 0.02	0.77 ± 0.03 ab	1.08 ± 0.01 b	1.31 ± 0.01
Wild taro	0.42 ± 0.02	0.75 ± 0.05 ab	1.18 ± 0.01 a	1.33 ± 0.01
Rodent tuber	0.41 ± 0.02	0.72 ± 0.02 b	1.07 ± 0.03 b	1.29 ± 0.03
Ornamental caladium	0.39 ± 0.01	0.80 ± 0.01 ab	1.14 ± 0.03 ab	1.30 ± 0.02
Lady finger banana	0.39 ± 0.01	0.84 ± 0.04 a	1.16 ± 0.01 ab	1.34 ± 0.02

^{ns}= not significant; Numbers followed by the same letter are not significantly different according to LSD 5%.

In this experiment we used aphid colony built up from single female, so genetical factor was assumed to have less influence to the aphid colony members.

The length of the antennae of *P. nigronervosa* was measured by using Milimeter v2.3.0 developed by Vis Tech. Project LLC after calibration. The antenna was removed from the aphid and put on the screen and bound the length to see the result of measurement. The aphid antennae varied only in the second and third instar, but there was no significant difference of antennal size in the first and fourth instar. Aphid antennal length correlate with the body length, shorter or longer than their body length, and *P. nigronervosa* had been reported to have antennae longer than their body length (Bagariang et al., 2019). This results of the aphid antennal measurement confirmed that the length of antennae of *P. nigronervosa* living in all experimental host plants were longer than their body length (Table 3). The measurement of body length and head width of the aphid was in the range of the measurement results reported by Bagariang et al. (2019).

The length of antennae and body length of *P. nigronervosa* lived on Bogor taro and rodent tuber was shorter than that of the aphid lived on banana, confirming the correlation between antennal length and body length of aphid species. The function of aphid antennae is also related to the safety of the aphid. The antennal sensors response to chemical and visual signal by inducing the formation of wings so the aphid has ability to avoid dangerous enemies or unfavorable conditions (Hu et al., 2019).

The morphometric of *P. nigronervosa* living on various Araceous plants species was significantly different between-some species (Table 1, 2 and 3), but the other morphological characteristics such body shape was relatively the same (Figure 1). Based on visual observation during the aphid development, the color of the aphid bodies was getting darker, from reddish brown in the first instar to blackish brown in the fourth instar. Host species did not affect the color of banana aphid. The color of the first instar was slightly transparent reddish brown, turned to greenish brown in the next two instars, getting darker in the fourth instar, and shiny black when reached the adult stage. Aphid color can be species-specific and is not influenced by nutritional quality of the host. According to Tsuchida (2016), aphid color is influenced by environment, especially the presence of natural enemies which may trigger physiological response in the form of color polymorphisms. In this experiment, the color of banana aphid used in the experiment was the same as that of the aphid in the rearing cage, because there was no natural

enemy in both sites.

The body shape development from the first to the fourth instar also similar amongst the aphid nymphs feeding on different hosts. This was not in accordance to the report by Mehrparvar et al. (2012) who found different morphology of black legume aphid (*Aphis craccivora*) living on different hosts, and Mdellel & Kamel (2015) who found the same phenomenon on *Pterochloroides persicae* living on different hosts. The first instar of *P. nigronervosa* was slightly angular in shape and becoming more oval along with the nymphal development to second, third and fourth instar. The segmentation of the antennae also noticeable from instar to instar. The first instar antennae consisted of 4 segments, the second and third instar had 5 segment antennae, and the fourth instar had the full length of six segment antennae (Figure 2). The length and segmentation of the aphid antennae were in accordance to the measurement of the aphid antennae reported by Bhadra & Agarwala (2012) that the length of *P. nigronervosa* antennae was about 1.01 to $1.35 \times$ its body length.

Biological Characteristics. Biological characteristics measured in the experiment were instar period, adult longevity, life span, and fecundity. Instar periods almost similar for each instar and there was no significant difference amongst host species (Table 4). The shortest instar period was 2.0 days, found in the second instar of the aphid living on wild taro and rodent tuber, while the longest instar period was 4.20 days found in the fourth instar of aphid nymph living on banana. The ranges of the four instar periods of *P. nigronervosa* live on banana were in accordance with the report by Padmanaban (2018) that *P. nigronervosa* had four nymphal instars and each instar was completed in 2-4 days. In all experimental hosts, nymphal period of the banana aphid was always longer in the fourth instar, while in the three previous instars the average instar periods almost the same, except that of Japanese taro which had second and third instar periods longer than those of other Araceous hosts. It is also noticeable that nymphal instar periods of *P. nigronervosa* living on banana was always significantly longer than the period of the aphid living on Araceous host. The difference of instar period between banana aphids living on banana and those living on Araceous plant might be in relation with nutritional quality of the hosts, where banana has better quality of nutrition and more suitable to the need of banana aphid. Both nymph and adult of *P. nigronervosa* required similar nutrition, and good quality of nutrition contained in their host would support nymphal growth better

than the hosts with less nutritional quality (Awmack & Leather, 2002).

Adult females started to reproduce parthenogenetically a day after the last molting of the fourth instar and keep reproducing for the rest of their life. *P. nigronervosa* actively gave birth to their offspring. The reproduction period differed significantly among host plant species, ranging from 12 to 38 days with an average of 24 days. The longest reproduction period was found in Bogor taro with an average of 38.3 days, while the shortest was found in Rodent tuber with

an average of 12.6 days (Table 5). The difference in the reproduction period had resulted in the difference in the number of offspring produced by adult females who lived on different host plants. The process of giving birth to the first instar nymph by an adult female was very slow and only one nymph was borne in each delivery. One adult female of *P. nigronervosa* was reported to produce 33.4 nymphs during its adult life (Basak et al., 2015). However, this research found that, on Araceous plants, an adult female of *P. nigronervosa* could give birth to a wider range of progenies, from 19.60 to 68.30




Figure 1. Morphological development of *P. nigronervosa* nymphs living on Araceous hosts and banana. A. Bogor taro; B. Belitung taro; C. Pontianak taro; D. Japanese taro; E. Wild taro; F. Rodent tuber; G. Ornamental caladium; H. Banana.




Figure 2. The development of the antennae *Pentalonia nigronervosa*. A. First instar; B. Second instar; C. Third instar; D. Fourth instar.

nymphs per adult female. The smallest fecundity was found in Rodent tuber, while the highest was found in Bogor taro (Table 5). Rodent tuber was reported to be one of the preferable alternative hosts of *P. nigronervosa* (Suparman et al., 2017), even though the fecundity of the aphid in this host was relatively low. Bogor taro was more preferable alternative host to *P. nigronervosa*, and its fecundity on this host was the highest. Host preference seems likely to relate to food quality contained in the host (Clissold & Simpson, 2015). Instead of having the highest fecundity, *P. nigronervosa* living on Bogor taro also had the longer average life span (47.9 days) and even longer than the aphid life span on banana plant as the main host, where the aphid had life span only 44.6 days (Table 5). All Araceous species used in this research were suitable as alternative host for *P. nigronervosa*, indicated by the aphid ability to feed on the plant and got nutrition to support their life and reproduction. Some species such as Bogor taro and Belitung taro could support *P. nigronervosa* even better than banana plant.

The average reproductive period in the experiment ranged from 13.10 to 38.30 (Table 5) and was wider than that reported by Basak et al. (2015) who reported

that reproductive period of *P. nigronervosa* in winter time ranged from 20.4 to 30.0 days. This difference could be caused by different temperature, because in this experiment the temperature was set at 25 °C, the temperature at which the aphid was at its best performance on adult longevity and fecundity. At the same temperature, the aphid was reported to reached its maximum efficiency in transmitting BBTv with incubation period about 35.32 days. Lower temperature reduced life span of the aphid and reduced transmission efficiency with longer incubation period, up to 189 days (Selvarajan & Balasubramanian 2013; Chakraborty et al., 2021).

The averages life span of *P. nigronervosa* living on Araceous plants ranged from 22.80 to 47.90 days, and the fecundity ranged from 26.40 to 68.30 nymphs per female (Table 5). The results were in wider ranges compared to those reported by Padmanaban (2018) that the life span of the aphid ranged from 27 to 37 days and the fecundity ranged from 35 to 50 nymphs per female. This difference could be caused by different temperature or rearing method as suggested by Xie et al. (2020). Data on the table Table 5 also show that

Table 4. Nymph instar period of *Pentalonia nigronervosa* lived on various host plants

Host Species	Instar period (days)			
	First instar	Second instar	Third instar	Fourth instar
Bogor Taro	2.20 ± 0.12 b	2.10 ± 0.09 b	2.20 ± 0.12 b	3.10 ± 0.09 bc
Belitung Taro	2.10 ± 0.09 b	2.10 ± 0.09 b	2.30 ± 0.14 b	3.50 ± 0.15 b
Pontianak Taro	2.30 ± 0.14 b	2.10 ± 0.09 b	2.50 ± 0.15 b	3.00 ± 0.20 bc
Japanese Taro	2.30 ± 0.14 b	2.50 ± 0.15 a	2.90 ± 0.09 a	3.40 ± 0.15 b
Wild Taro	2.10 ± 0.09 b	2.00 ± 0.00 b	2.30 ± 0.14 b	3.40 ± 0.20 b
Rodent Tuber	2.20 ± 0.12 b	2.00 ± 0.00 b	2.40 ± 0.15 b	3.60 ± 0.20 b
Ornamental Caladium	2.30 ± 0.14 b	2.50 ± 0.15 a	2.10 ± 0.00 b	2.70 ± 0.14 c
Lady Finger Banana	3.00 ± 0.00 a	2.80 ± 0.12 a	3.00 ± 0.00 a	4.20 ± 0.25 a

Number followed by the same letter are not significantly different according to LSD 5%.

Table 5. The reproductive period of *Pentalonia nigronervosa* on various host plants

Host species	The reproductive period (days)	Life span (days)	Fecundity (nymph/female)
Bogor Taro	38.30 ± 3.64 a	47.90 ± 3.75 a	68.30 ± 7.78 a
Belitung Taro	26.30 ± 2.80 b	36.50 ± 2.94 b	37.70 ± 3.49 bc
Pontianak Taro	23.60 ± 1.79 b	33.50 ± 1.83 b	40.10 ± 4.60 bc
Japanese Taro	13.10 ± 1.30 c	23.20 ± 1.10 b	26.40 ± 3.45 cd
Wild Taro	23.90 ± 1.85 b	33.70 ± 1.78 b	33.60 ± 3.44 c
Rodent Tuber	12.60 ± 1.53 c	22.80 ± 1.46 b	19.60 ± 1.34 d
Ornamental Caladium	21.90 ± 1.17 b	31.20 ± 1.16 b	40.10 ± 3.92 bc
Lady Finger Banana	32.70 ± 0.63 ab	44.60 ± 1.45 a	51.50 ± 1.36 b

Numbers followed by the same letters are not significantly different according to LSD 5%.

the fecundity of *P. nigronervosa* living on banana had significantly higher than the aphid living on Araceous plants, except Bogor taro where the aphid had the highest fecundity, even significantly higher than banana itself. The biological performance of banana aphid on Bogor taro was outstanding with longer reproductive period, longer life span and higher fecundity. This is an indication that banana and Bogor taro had better host quality for the aphid, and the aphid itself is able to live on good as well as on poor-quality hosts. Banana as the main host of *P. nigronervosa* has had long good interaction with the aphid, indicating that the interaction is supported by good quality of nutrition provided by the host. A poor-quality host plant would cause a female insect produce only few good-quality progenies or a large number of poor-quality progenies (Gilkeson, 2012). In this experiment, we used young suckers to facilitate the aphid with better nutrient quality, since nymphal nutrition intake had strong influence to the aphid fecundity. The main food of aphids is phloem sap with high content of sugar and low content of amino acids (Shaaban et al., 2020) and to obtain enough amino acid, aphids suck a lot of phloem sap and secrete the sugar in the form of honey dew. The amino acid content of phloem sap was also very important for aphid to infest new plant host (Liu et al., 2016).

Even though many insects had considerable plasticity to their host plant quality (Awmack and Leather, 2002), the quality of the host might affect the quality of the insect progenies but not the quantity. In this research, the progenies were removed from the reproducing female so the quality of the progenies was not intensively studied. However, the aphid had been reared in Araceous plants prior to its used in the biological assay experiment, and the aphid could produce numbers of progenies needed by the experiment. Furthermore, we prefer to measure fecundity of the aphid without considering the viability of the progenies which determine their fertility. The difference between fecundity and fertility of *P. nigronervosa* in relation to host quality was clearly reported by Safari et al. (2022) that banana plants infected by BBTV reduced fertility of the aphid but increases its fecundity.

CONCLUSION

Banana aphid *P. nigronervosa* was able to live, feed and reproduce on Araceous plants with most morphometric and biological characteristics were not significantly different from those of the aphids living on banana. However, the aphids living on Bogor taro showed significantly longer life span and

higher fecundity compared to those living on other hosts, including banana. Since Araceous plants were abundantly available surrounding banana cultivation, the interaction between banana aphid and Araceous plants would have implication on BBTVD epidemiology. The long persistence of BBTVD in the body of viruliferous aphid and the tendency of the aphid to leave infected banana will encourage the viruliferous aphid to migrate to Araceous plants which have ability to reduce the transmission efficiency of BBTVD by the aphid.

ACKNOWLEDGMENTS

We would like to thank Rector of Universitas Sriwijaya who have financially supported this research through the scheme of Excellent Competitive Research Program, and the Dean of Faculty of Agriculture Universitas Sriwijaya for the free used of laboratories and experimental garden during the research.

FUNDING

The research behind this paper was a part of research entitled “Effects of various species of the alternative host of *Banana bunchy top virus* vector on its transmission efficiency and the virus pathogenicity” funded by Universitas Sriwijaya through Excellent Competitive Research Funding, 2021.

AUTHORS' CONTRIBUTIONS

SS was the main contributor of this manuscript. SS and HH concepted and designed the experiment. RO and AA conducted survey for banana aphid collection. All authors involved in the laboratory and field experiments. SS prepared the manuscript, and all authors read and approved the final manuscript.

COMPETING INTERESTS

The Authors declare that there is no competing interest in the publication of this manuscript.

REFERENCES

- Aqueel MA, Collins CM, Raza AM, Ahmad S, Tariq M, & Leather SR. 2013. Effect of plant nutrition on aphid size, prey consumption, and life history characteristics of green lacewing. *Insect Sci.* 21(1): 74–82. <https://doi.org/10.1111/1744-7917.12019>

- Awmack CS & Leather SR. 2002. Host plant quality and fecundity in herbivorous insects. *Annu. Rev. Entomol.* 47: 817–44. <https://doi.org/10.1146/annurev.ento.47.091201.145300>
- Bagariang W, Hidayat P, & Hidayat SH. 2019. Morphometric analysis and host range of the genus *Pentalonia* Coquerel (Hemiptera: Aphididae) infesting banana in Java. *Jurnal Perlindungan Tanaman Indonesia.* 23(2): 171–178. <https://doi.org/10.22146/jpti.38220>
- Basak G, Banerjee A, & Bandyopadhyay B. 2015. Studies on some bio-ecological aspects and varietal preference of banana aphid, *Pentalonia nigronervosa* Coquerel (Hemiptera: Aphididae). *J. Crop Weed.* 11(2): 181–186.
- Bhadra P & Agarwala BK. 2012. On the morphological and genotypic variations of two congeneric species of banana aphid *Pentalonia* (Homoptera: Aphididae) from India. *Adv. Life Sci.* 2(3): 75–81. <https://doi.org/10.5923/j.als.20120203.06>
- Biale H, Mendel Z, & Soroker V. 2017. Insects associated with the banana aphid *Pentalonia nigronervosa* Coquerel (Hemiptera: Aphididae) in banana plantations with special emphasis on the ant community. *Phytoparasitica* 45: 361–372. <https://doi.org/10.1007/s12600-017-0592-z>
- Bragard C, Caciagli P, Lemaire O, Lopez-Moya JJ, MacFarlane S, Peters D, Susi P, & Torrance L. 2013. Status and prospects of plant virus control through interference with vector transmission. *Annu. Rev. Phytopathol.* 51: 177–201. <https://doi.org/10.1146/annurev-phyto-082712-102346>
- Chakraborty S, Barman M, Samanta S, Roy M, & Tarafdar J. 2021. Effect of banana bunchy top virus on the heat shock protein genes of *Pentalonia nigronervosa* during temperature susceptibility and its effect on virus transmission. *Agronomy* 11(9): 1866. <https://doi.org/10.3390/agronomy11091866>
- Chen Y & Hu X. 2013. High-throughput detection of *Banana bunchy top virus* in banana plants and aphids using real-time TaqMan® PCR. *J. Virol. Methods.* 193(1): 177–183. <https://doi.org/10.1016/j.jviromet.2013.06.013>
- Clissold FJ & Simpson SJ. 2015. Temperature, food quality and life history traits of herbivorous insects. *Curr. Opin. Insect Sci.* 11: 63–70. <https://doi.org/10.1016/j.cois.2015.10.011>
- Duay JAM, Miller RH, Wall GC, Pike KS, & Footitt RG. 2014. *Pentalonia nigronervosa* Coquerel and *Pentalonia caladii* van der Goot (Hemiptera: Aphididae) and their relationship to *Banana bunchy top virus* in Micronesia. *Pac. Sci.* 68(3): 359–364. <https://doi.org/10.2984/68.3.5>
- Elayabalan S, Subramaniam S, & Selvarajan S. 2015. Banana bunchy top disease (BBTD) symptom expression in banana and strategies for transgenic resistance: A review. *Emir. J. Food Agric.* 27(1): 55–74. <https://doi.org/10.9755/ejfa.v27i1.19197>
- Footitt RG, Maw HEL, Pike KS, & Miller RH. 2010. The identity of *Pentalonia nigronervosa* Coquerel and *P. caladii* van der Got (Hemiptera: Aphididae) based on molecular and morphometric analysis. *Zootaxa* 2358: 25–38.
- Footitt RG & Maw HEL. 2019. Geographic distribution, host preferences and molecular diversity within the genus *Pentalonia* (Hemiptera: Aphididae). *Zootaxa* 4701(4): 383–391. <https://doi.org/10.11646/zootaxa.4701.4.4>
- Gao Y, Ren R, Peng J, Wang D, Shi X, Zheng L, Zhang Z, Zhu C, Liu Y, Dai L, & Zhang D. 2021. The *Gustavus* gene can regulate the fecundity of the green peach aphid, *Myzus persicae* (Sulzer). *Front. Physiol.* 11: 596392. <https://doi.org/10.3389/fphys.2020.596392>
- Gilkeson LA. 2012. A note on fecundity of the aphid predator, *Aphidoletes aphidimyza* (Rondani) (Diptera: Cecidomyiidae) *Can. Entomol.* 119 (12):1145–1146. <https://doi.org/10.4039/Ent1191145-12>
- Hu L, Gui W, Chen B, & Chen L. 2019. Transcriptome profiling of maternal stress-induced wing dimorphism in pea aphids. *Ecol. Evol.* 9(20): 11848–11862. <https://doi.org/10.1002/ece3.5692>
- Jebakumar RM, Balasubramanian V, & Selvarajan R. 2018. Virus titre determines the efficiency of *Pentalonia nigronervosa* (Aphididae: Hemiptera) to transmit *Banana bunchy top virus*. *VirusDis.* 29(4): 499–505. <https://doi.org/10.1007/s13337-018-0493-x>
- Jekayinoluwa T, Tripathi L, Tripathi JN, Ntui VO, Obiero G, Muge E, & Dale J. 2020. RNAi technology for management of banana bunchy top disease. *Food Energy Secur.* 9(4): e247. <https://doi.org/10.1002/fes3.247>
- Jekayinoluwa T, Tripathi JN, Dugdale B, Obiero G,

- Muge E, Dale J, & Tripathi L. 2021. Transgenic expression of dsRNA targeting the *Pentalonia nigronervosa* acetylcholinesterase gene in banana and plantain reduces aphid populations. *Plants* 10(4): 613. <https://doi.org/10.3390/plants10040613>
- Le Guigo P, Rolier A, & Le Corff J. 2012. Plant neighborhood influences colonization of Brassicaceae by specialist and generalist aphids. *Oecologia* 169(3): 753–761. <https://doi.org/10.1007/s00442-011-2241-4>
- Liu YH, Kang ZW, Guo Y, Zhu GS, Shah MMR, Song Y, Fan YL, Jing X, & Liu TX. 2016. Nitrogen hurdle of host alternation for a polyphagous aphid and the associated changes of endosymbionts. *Sci. Rep.* 6: 24781. <https://doi.org/10.1038/srep24781>
- Mdellel L & Kamel MBH. 2015. Effect of host plant on morphology of *Pterochloroides persicae* Cholodkovsky 1899 (Hemiptera, Aphididae). *J. Entomol. Zool. Stud.* 3(3): 324–327.
- Mehrpour M, Madjdzadeh SM, Mahdavi Arab N, Esmailbeygi M, & Ebrahimpour E. 2012. Morphometric discrimination of Black Legume Aphid, *Aphis craccivora* Koch (Hemiptera: Aphididae), populations associated with different host plants. *North-West. J. Zool.* 8(1): 172–180.
- Niyongere C, Losenge T, Ateka EM, Ntukamazina N, Ndayiragije P, Simbare A, Cimpaye P, Nintije P, Lepoint P, & Blomme G. 2013. Understanding banana bunchy top disease epidemiology in Burundi for an enhanced and integrated management approach. *Plant Pathol.* 62(3): 562–570. <https://doi.org/10.1111/j.1365-3059.2012.02676.x>
- Oktarida R, Suparman S, & Hamidson H. 2022. Pengaruh spesies tumbuhan Araceae terhadap efisiensi penularan *Banana bunchy top virus* oleh *Pentalonia nigronervosa* [The effects of Araceous plant species on the efficiency of *Banana bunchy top virus* transmission by *Pentalonia nigronervosa*]. *Sainmatika: Jurnal Ilmiah Matematika dan Ilmu Pengetahuan Alam.* 19(1): 66–73. <https://doi.org/10.31851/sainmatika.v19i1.8094>
- Padmanaban B. 2018. Pest of banana. In: Omcar (ed). *Pest and their management*. pp. 441–455. Springer. Singapore.
- Qazi J. 2016. *Banana bunchy top virus* and the bunchy top disease. *J. Gen. Plant Pathol.* 82(1): 2–11. <https://doi.org/10.1007/s10327-015-0642-7>
- Martínez RAF & Costamagna AC. 2018. Effects of crowding and host plant quality on morph determination in the soybean aphid, *Aphis glycines*. *Entomol. Exp. Appl.* 166(1): 53–62. <https://doi.org/10.1111/eea.12637>
- Safari IM, Tougeron K, Bragard C, Basengere EB, Masamba JW, & Hance T. 2022. Trade-off between longevity, body length and fecundity in *Pentalonia nigronervosa* after *Banana bunchy top virus* infection. *Conference on Biodiversity in a rapidly changing world*. Louvain-la-Neuve, Belgium April 2022.
- Selvarajan R & Balasubramanian V. 2013. Natural Occurrence of *Banana bunchy top virus* in *Ensete superbum* in India. *Indian J. Virol.* 24(1): 97–98. <https://doi.org/10.1007/s13337-012-0123-y>
- Shaaban B, Seeburger V, Schroeder A, & Lohaus G. 2020. Sugar, amino acid and inorganic ion profiling of the honeydew from different hemipteran species feeding on *Abies alba* and *Picea abies*. *PLoS ONE* 15(1): e0228171. <https://doi.org/10.1371/journal.pone.0228171>
- Suparman, Gunawan B, Pujiastuti Y, Arsi, & Cameron RR. 2017. Alternative hosts of banana aphid *Pentalonia nigronervosa* Coq. (Hemiptera: Aphididae), the vector transmitting Banana Bunchy Top Virus. *J. Adv. Agric. Technol.* 4(4): 354–359. <https://doi.org/10.18178/joaat.4.4.354-359>
- Tricahyati T, Suparman S, & Irsan C. 2022. Pengaruh umur kematian tanaman sumber inokulum *Banana bunchy top virus* terhadap efisiensi penularannya [Effect of time mortality of plant which was become an inoculum source of *Banana bunchy top virus* on its transmission efficiency]. *Sainmatika: Jurnal Ilmiah Matematika dan Ilmu Pengetahuan Alam.* 19(1): 55–65. <https://doi.org/10.31851/sainmatika.v19i1.7977>
- Tsuchida T. 2016. Molecular basis and ecological relevance of aphid body colors. *Curr. Opin. Insect Sci.* 17: 74–80. <https://doi.org/10.1016/j.cois.2016.07.005>
- Watanabe S & Bressan A. 2013. Tropism, compartmentalization and retention of *Banana bunchy top virus* (Nanoviridae) in the aphid vector *Pentalonia nigronervosa*. *J. Gen. Virol.* 94(1):

- 209–219. <https://doi.org/10.1099/vir.0.047308-0>
- Watanabe S, Greenwell AM, & Bressan A. 2013. Localization, concentration, and transmission efficiency of *Banana bunchy top virus* in four asexual lineages of *Pentalonia aphids*. *Viruses* 5(2): 758–776. <https://doi.org/10.3390/v5020758>
- Xie H, Shi J, Shi F, Xu H, He K, & Wang Z. 2020. Aphid fecundity and defenses in wheat exposed to a combination of heat and drought stress. *J. Exp. Bot.* 71(9): 2713–2722. <https://doi.org/10.1093/jxb/eraa017>